

Intrinsic geometric structures of quantum integrable systems

Anton M. Zeitlin

Georgia Institute of Technology

University of Virginia

Charlottesville

March 5, 2025

Example of integrable system: harmonic oscillator

Anton M. Zeitlin

2-dimensional **phase space** with coordinate q and momentum p :

Hamiltonian:
$$H = \frac{p^2 + q^2}{2},$$

Poisson bracket:
$$\{F, G\} = \frac{\partial F}{\partial p} \frac{\partial G}{\partial q} - \frac{\partial G}{\partial p} \frac{\partial F}{\partial q}.$$

Equations of motion:
$$\left. \begin{aligned} \frac{dq}{dt} &= \{H, q\} = p \\ \frac{dp}{dt} &= \{H, p\} = -q \end{aligned} \right\} \Rightarrow \frac{d^2 q}{dt^2} + q = 0$$

Action-angle variables: polar coordinates in (q, p) -space

Energy level set: $L_E = \{p^2 + q^2 = 2E\}.$

Equations of motion for action-angle variables (H, ϕ) :

$$\frac{d\phi}{dt} = 1, \quad \frac{dH}{dt} = 0$$

Introduction

Quantum Integrable
models

Enumerative geometry
and Bethe ansatz

QQ-systems and
(G, \hbar)-opers

Applications

Example of integrable system: harmonic oscillator

Anton M. Zeitlin

2-dimensional **phase space** with coordinate q and momentum p :

Hamiltonian:
$$H = \frac{p^2 + q^2}{2},$$

Poisson bracket:
$$\{F, G\} = \frac{\partial F}{\partial p} \frac{\partial G}{\partial q} - \frac{\partial G}{\partial p} \frac{\partial F}{\partial q}.$$

Equations of motion:
$$\left. \begin{aligned} \frac{dq}{dt} &= \{H, q\} = p \\ \frac{dp}{dt} &= \{H, p\} = -q \end{aligned} \right\} \Rightarrow \frac{d^2 q}{dt^2} + q = 0$$

Action-angle variables: polar coordinates in (q, p) -space

Energy level set: $L_E = \{p^2 + q^2 = 2E\}.$

Equations of motion for action-angle variables (H, ϕ) :

$$\frac{d\phi}{dt} = 1, \quad \frac{dH}{dt} = 0$$

Introduction

Quantum Integrable
models

Enumerative geometry
and Bethe ansatz

QQ-systems and
 (G, \hbar) -opers

Applications

Example of integrable system: harmonic oscillator

Anton M. Zeitlin

2-dimensional **phase space** with coordinate q and momentum p :

Hamiltonian:
$$H = \frac{p^2 + q^2}{2},$$

Poisson bracket:
$$\{F, G\} = \frac{\partial F}{\partial p} \frac{\partial G}{\partial q} - \frac{\partial G}{\partial p} \frac{\partial F}{\partial q}.$$

Equations of motion:
$$\left. \begin{aligned} \frac{dq}{dt} &= \{H, q\} = p \\ \frac{dp}{dt} &= \{H, p\} = -q \end{aligned} \right\} \Rightarrow \frac{d^2 q}{dt^2} + q = 0$$

Action-angle variables: polar coordinates in (q, p) -space

Energy level set: $L_E = \{p^2 + q^2 = 2E\}.$

Equations of motion for action-angle variables (H, ϕ) :

$$\frac{d\phi}{dt} = 1, \quad \frac{dH}{dt} = 0$$

Introduction

Quantum Integrable
models

Enumerative geometry
and Bethe ansatz

QQ-systems and
(G, \hbar)-opers

Applications

Classical integrable systems: what are they?

Anton M. Zeitlin

Symplectic $2n$ -manifold M : phase space, which has information of coordinates and momenta of a physical system.

Equations of motion:

$$\frac{df}{dt} = \{H, f\}.$$

Integrability: family of conserved quantities: $\{F_i\}_{i=1}^n$:

$$\{F_i, F_j\} = 0, \quad F_1 = H.$$

Liouville-Arnold theorem:

- ▶ Compact connected components of $L_c = \{F_i = c_i\}_{i=1}^n$ are diffeomorphic to \mathbb{T}^n .
- ▶ Existence of action-angle variables $\{l_i\}_{i=1}^n, \{\phi^i\}_{i=1}^n$ in the neighborhood of L_c :

$$\frac{d\phi^i}{dt} = \omega^i, \quad \frac{dl_i}{dt} = 0.$$

Finding action/angle variables is a non-trivial problem.

Introduction

Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

Classical integrable systems: what are they?

Anton M. Zeitlin

Symplectic $2n$ -manifold M : phase space, which has information of coordinates and momenta of a physical system.

Equations of motion:

$$\frac{df}{dt} = \{H, f\}.$$

Integrability: family of conserved quantities: $\{F_i\}_{i=1}^n$:

$$\{F_i, F_j\} = 0, \quad F_1 = H.$$

Liouville-Arnold theorem:

- ▶ Compact connected components of $L_c = \{F_i = c_i\}_{i=1}^n$ are diffeomorphic to \mathbb{T}^n .
- ▶ Existence of action-angle variables $\{I_i\}_{i=1}^n, \{\phi^i\}_{i=1}^n$ in the neighborhood of L_c :

$$\frac{d\phi^i}{dt} = \omega^i, \quad \frac{dI_i}{dt} = 0.$$

Finding action/angle variables is a non-trivial problem.

Introduction

Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

Classical integrable systems: what are they?

Anton M. Zeitlin

Symplectic $2n$ -manifold M : phase space, which has information of coordinates and momenta of a physical system.

Equations of motion:

$$\frac{df}{dt} = \{H, f\}.$$

Integrability: family of conserved quantities: $\{F_i\}_{i=1}^n$:

$$\{F_i, F_j\} = 0, \quad F_1 = H.$$

Liouville-Arnold theorem:

- ▶ Compact connected components of $L_c = \{F_i = c_i\}_{i=1}^n$ are diffeomorphic to \mathbb{T}^n .
- ▶ Existence of action-angle variables $\{I_i\}_{i=1}^n, \{\phi^i\}_{i=1}^n$ in the neighborhood of L_c :

$$\frac{d\phi^i}{dt} = \omega^i, \quad \frac{dI_i}{dt} = 0.$$

Finding action/angle variables is a non-trivial problem.

Introduction

Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

Classical integrable systems: what are they?

Anton M. Zeitlin

Symplectic $2n$ -manifold M : phase space, which has information of coordinates and momenta of a physical system.

Equations of motion:

$$\frac{df}{dt} = \{H, f\}.$$

Integrability: family of conserved quantities: $\{F_i\}_{i=1}^n$:

$$\{F_i, F_j\} = 0, \quad F_1 = H.$$

Liouville-Arnold theorem:

- ▶ Compact connected components of $L_c = \{F_i = c_i\}_{i=1}^n$ are diffeomorphic to \mathbb{T}^n .
- ▶ Existence of action-angle variables $\{I_i\}_{i=1}^n, \{\phi^i\}_{i=1}^n$ in the neighborhood of L_c :

$$\frac{d\phi^i}{dt} = \omega^i, \quad \frac{dI_i}{dt} = 0.$$

Finding action/angle variables is a non-trivial problem.

Introduction

Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

Explosion of interest in integrable systems: 60s -70s

Anton M. Zeitlin

Integrable soliton equations in (1+1)-dimensions,
e.g. Korteweg-de Vries (KdV) equation:

$$u_t = -u_{xxx} + 6uu_x.$$

C. S. Gardner, J. M. Greene, M. D. Kruskal, R. Miura'67; P. Lax'68;

L. Faddeev, V. Zakharov'71

Lie-theoretic methods through Lax pair formulation:

$$\frac{dL}{dt} = [A, L],$$

where $L = -\partial_x^2 + u(x, t)$ for KdV.

I. Gelfand, L. Dickey'76; V. Drinfeld, V. Sokolov'85

Inverse Scattering Method (ISM):

spectral data of $L \rightarrow$ action-angle variables

At the same time many finite-dimensional multiparticle integrable systems were discovered: Calogero-Moser, Toda, Ruijsenaars-Schneider, etc.

Introduction

Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

Explosion of interest in integrable systems: 60s -70s

Anton M. Zeitlin

Integrable soliton equations in (1+1)-dimensions,
e.g. Korteweg-de Vries (KdV) equation:

$$u_t = -u_{xxx} + 6uu_x.$$

C. S. Gardner, J. M. Greene, M. D. Kruskal, R. Miura'67; P. Lax'68;

L. Faddeev, V. Zakharov'71

Lie-theoretic methods through Lax pair formulation:

$$\frac{dL}{dt} = [A, L],$$

where $L = -\partial_x^2 + u(x, t)$ for KdV.

I. Gelfand, L. Dickey'76; V. Drinfeld, V. Sokolov'85

Inverse Scattering Method (ISM):

spectral data of $L \rightarrow$ action-angle variables

At the same time many finite-dimensional multiparticle integrable systems were discovered: Calogero-Moser, Toda, Ruijsenaars-Schneider, etc.

Introduction

Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

Explosion of interest in integrable systems: 60s -70s

Anton M. Zeitlin

Integrable soliton equations in (1+1)-dimensions,
e.g. Korteweg-de Vries (KdV) equation:

$$u_t = -u_{xxx} + 6uu_x.$$

C. S. Gardner, J. M. Greene, M. D. Kruskal, R. Miura'67; P. Lax'68;

L. Faddeev, V. Zakharov'71

Lie-theoretic methods through Lax pair formulation:

$$\frac{dL}{dt} = [A, L],$$

where $L = -\partial_x^2 + u(x, t)$ for KdV.

I. Gelfand, L. Dickey'76; V. Drinfeld, V. Sokolov'85

Inverse Scattering Method (ISM):

spectral data of $L \rightarrow$ action-angle variables

At the same time many finite-dimensional multiparticle integrable systems were discovered: Calogero-Moser, Toda, Ruijsenaars-Schneider, etc.

Introduction

Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

Integrable soliton equations in (1+1)-dimensions,
e.g. Korteweg-de Vries (KdV) equation:

$$u_t = -u_{xxx} + 6uu_x.$$

C. S. Gardner, J. M. Greene, M. D. Kruskal, R. Miura'67; P. Lax'68;

L. Faddeev, V. Zakharov'71

Lie-theoretic methods through Lax pair formulation:

$$\frac{dL}{dt} = [A, L],$$

where $L = -\partial_x^2 + u(x, t)$ for KdV.

I. Gelfand, L. Dickey'76; V. Drinfeld, V. Sokolov'85

Inverse Scattering Method (ISM):

spectral data of $L \rightarrow$ action-angle variables

At the same time many finite-dimensional multiparticle integrable systems were discovered: Calogero-Moser, Toda, Ruijsenaars-Schneider, etc.

Introduction

Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

Quantum integrability:

$$[H_i, H_j] = 0, \quad H_i : \mathcal{H} \rightarrow \mathcal{H}$$

Finding action/angle variables \rightarrow simultaneous diagonalization of H_i .

Quantization of (1+1)-models? Put them on the lattice.

Lattice integrable models \rightarrow new algebraic structures:

R-matrix and Yang-Baxter equation

accompanied with

algebraic Bethe ansatz

lead to the the discovery of Quantum inverse scattering method (QISM)
developed by Leningrad School.

That eventually led to the discovery of quantum groups by Drinfeld and Jimbo.

Introduction

Quantum Integrable
models

Enumerative geometry
and Bethe ansatz

QQ-systems and
(G, \hbar)-opers

Applications

Quantum integrability:

$$[H_i, H_j] = 0, \quad H_i : \mathcal{H} \rightarrow \mathcal{H}$$

Finding action/angle variables \rightarrow simultaneous diagonalization of H_i .

Quantization of (1+1)-models? Put them on the lattice.

Lattice integrable models \rightarrow new algebraic structures:

R-matrix and **Yang-Baxter equation**

accompanied with

algebraic Bethe ansatz

lead to the the discovery of **Quantum inverse scattering method (QISM)**
developed by **Leningrad School**.

That eventually led to the discovery of **quantum groups** by **Drinfeld** and **Jimbo**.

Introduction

Quantum Integrable
models

Enumerative geometry
and Bethe ansatz

QQ-systems and
(G, \hbar)-opers

Applications

Quantum integrability:

$$[H_i, H_j] = 0, \quad H_i : \mathcal{H} \rightarrow \mathcal{H}$$

Finding action/angle variables \rightarrow simultaneous diagonalization of H_i .

Quantization of (1+1)-models? Put them on the lattice.

Lattice integrable models \rightarrow new algebraic structures:

R-matrix and **Yang-Baxter equation**

accompanied with

algebraic Bethe ansatz

lead to the the discovery of **Quantum inverse scattering method (QISM)**
developed by **Leningrad School**.

That eventually led to the discovery of **quantum groups** by **Drinfeld** and **Jimbo**.

Introduction

Quantum Integrable
models

Enumerative geometry
and Bethe ansatz

QQ-systems and
(G, \hbar)-opers

Applications

Quantum integrability:

$$[H_i, H_j] = 0, \quad H_i : \mathcal{H} \rightarrow \mathcal{H}$$

Finding action/angle variables \rightarrow simultaneous diagonalization of H_i .

Quantization of (1+1)-models? Put them on the lattice.

Lattice integrable models \rightarrow new algebraic structures:

R-matrix and **Yang-Baxter equation**

accompanied with

algebraic Bethe ansatz

lead to the the discovery of **Quantum inverse scattering method (QISM)**
developed by **Leningrad School**.

That eventually led to the discovery of **quantum groups** by **Drinfeld** and **Jimbo**.

Introduction

Quantum Integrable
models

Enumerative geometry
and Bethe ansatz


QQ-systems and
(G, \hbar)-opers

Applications

Introduction

Quantum Integrable
modelsEnumerative geometry
and Bethe ansatzQQ-systems and
(G, \hbar)-opers

Applications



R.P. Feynman: "I got really fascinated by these (1+1)-dimensional models that are solved by the Bethe ansatz and how mysteriously they jump out at you and work and you don't know why. I am trying to understand all this better."

Introduction

Quantum integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

- ▶ Dubrovin, Givental, Kontsevich, Witten established first relations with integrability in the context of enumerative geometry.

Notable cases:

- ▶ Witten's conjecture, proven by Kontsevich, relating intersection numbers on the moduli space of curves and the τ -function of KdV model.
- ▶ Givental and collaborators: description of the enumerative geometry of flag varieties (quantum cohomology/quantum K-theory) via classical and quantum multiparticle systems of Toda type.
- ▶ Feigin, Frenkel, and collaborators, while studying conformal field theory/representation theory of affine Lie algebras, discovered the relation:
Connections on \mathbb{P}^1 called **opers** \leftrightarrow **Gaudin integrable model**
That turned out to be an example of the **geometric Langlands correspondence**.

Introduction

Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

- ▶ Dubrovin, Givental, Kontsevich, Witten established first relations with integrability in the context of enumerative geometry.

Notable cases:

- ▶ Witten's conjecture, proven by Kontsevich, relating intersection numbers on the moduli space of curves and the τ -function of KdV model.
- ▶ Givental and collaborators: description of the enumerative geometry of flag varieties (quantum cohomology/quantum K-theory) via classical and quantum multiparticle systems of Toda type.
- ▶ Feigin, Frenkel, and collaborators, while studying conformal field theory/representation theory of affine Lie algebras, discovered the relation:

Connections on \mathbb{P}^1 called **opers** \leftrightarrow **Gaudin integrable model**

That turned out to be an example of the **geometric Langlands correspondence**.

Introduction

Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

► Nakajima, Schiffmann, Varagnolo-Vasserot:

Geometric realization of representations of quantum groups on cohomology and K-theory of **symplectic resolutions**, in particular, on **Nakajima quiver varieties**.

Okounkov:

“Symplectic resolutions are the Lie algebras of XXI century”

► 2010s: Nekrasov, Shatashvili:

Hints from supersymmetric gauge theory → geometric realization of quantum integrable models solved by Bethe ansatz.

Okounkov and his school: enumerative geometry of symplectic resolutions.

Introduction

Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

► Nakajima, Schiffmann, Varagnolo-Vasserot:

Geometric realization of representations of quantum groups on cohomology and K-theory of **symplectic resolutions**, in particular, on **Nakajima quiver varieties**.

Okounkov:

“Symplectic resolutions are the Lie algebras of XXI century”

► 2010s: Nekrasov, Shatashvili:

Hints from supersymmetric gauge theory → geometric realization of quantum integrable models solved by Bethe ansatz.

Okounkov and his school: enumerative geometry of symplectic resolutions.

Fusion of:

- ▶ Theory of integrable systems
- ▶ Geometric representation theory
- ▶ Enumerative geometry
- ▶ Supersymmetric gauge theories

More concretely, we will discuss the following:

- ▶ Nekrasov-Shatashvili conjectures:
Bethe ansatz solution for quantum integrable systems encodes enumerative invariants of certain symplectic resolutions:
quantum cohomology, quantum K-theory.
- ▶ On the other hand, geometrization of the relations in the corresponding rings lead to the deformation of the version of geometric Langlands correspondence by Feigin-Frenkel.
- ▶ Applications bring together many parts of theoretical physics and mathematics, such as quantum-classical duality, cluster algebras, and 3D mirror symmetry.

Introduction

Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

Fusion of:

- ▶ Theory of integrable systems
- ▶ Geometric representation theory
- ▶ Enumerative geometry
- ▶ Supersymmetric gauge theories

More concretely, we will discuss the following:

- ▶ **Nekrasov-Shatashvili** conjectures:
Bethe ansatz solution for quantum integrable systems encodes enumerative invariants of certain symplectic resolutions:
quantum cohomology, quantum K-theory.
- ▶ On the other hand, geometrization of the relations in the corresponding rings lead to the deformation of the version of geometric Langlands correspondence by Feigin-Frenkel.
- ▶ Applications bring together many parts of theoretical physics and mathematics, such as quantum-classical duality, cluster algebras, and 3D mirror symmetry.

Introduction

Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

Fusion of:

- ▶ Theory of integrable systems
- ▶ Geometric representation theory
- ▶ Enumerative geometry
- ▶ Supersymmetric gauge theories

More concretely, we will discuss the following:

- ▶ **Nekrasov-Shatashvili** conjectures:
Bethe ansatz solution for quantum integrable systems encodes enumerative invariants of certain symplectic resolutions:
quantum cohomology, quantum K-theory.
- ▶ On the other hand, geometrization of the relations in the corresponding rings lead to the deformation of the version of geometric Langlands correspondence by Feigin-Frenkel.
- ▶ Applications bring together many parts of theoretical physics and mathematics, such as quantum-classical duality, cluster algebras, and 3D mirror symmetry.

Introduction

Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

Fusion of:

- ▶ Theory of integrable systems
- ▶ Geometric representation theory
- ▶ Enumerative geometry
- ▶ Supersymmetric gauge theories

More concretely, we will discuss the following:

- ▶ **Nekrasov-Shatashvili** conjectures:
Bethe ansatz solution for quantum integrable systems encodes enumerative invariants of certain symplectic resolutions:
quantum cohomology, quantum K-theory.
- ▶ On the other hand, geometrization of the relations in the corresponding rings lead to the deformation of the version of geometric Langlands correspondence by Feigin-Frenkel.
- ▶ Applications bring together many parts of theoretical physics and mathematics, such as quantum-classical duality, cluster algebras, and 3D mirror symmetry.

Introduction

Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

P. Koroteev, P. Pushkar, E. Frenkel, D. Sage, A. Smirnov

- ▶ P. Koroteev, A.Z., *The Zoo of Opers and Dualities*,
Int. Math. Res. Not.'24
- ▶ P. Koroteev, E. Frenkel, D. Sage, A.Z., *q-Opers, QQ-Systems, and Bethe Ansatz*,
J. Eur. Math. Soc.'24
- ▶ P. Koroteev, A. Z., *3d Mirror Symmetry for Instanton Moduli Spaces*,
Comm. Math. Phys.'23
- ▶ P. Koroteev, A.Z., *q-Opers, QQ-Systems, and Bethe Ansatz II: Generalized Minors*,
J. Reine Angew. Math.'23
- ▶ P. Koroteev, A.Z., *Toroidal q-Opers*,
J. Inst. Math. Jussieu'23
- ▶ P. Koroteev, D. Sage, A.Z., *$(SL(N), q)$ -opers, the q -Langlands correspondence, and quantum/classical duality*,
Comm. Math. Phys.'21
- ▶ P. Koroteev, P. Pushkar, A. Smirnov, A.Z., *Quantum K-theory of Quiver Varieties and Many-Body Systems*,
Selecta Math. '21
- ▶ P. Koroteev, A.Z., *qKZ/tRS Duality via Quantum K-Theoretic Counts*,
Math. Res. Lett.'21
- ▶ P. Pushkar, A. Smirnov, A.Z., *Baxter Q-operator from quantum K-theory*,
Adv. Math.'20

Introduction

Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

Let us consider Lie algebra \mathfrak{g} .

The associated *loop algebra* is $\hat{\mathfrak{g}} = \mathfrak{g}[t, t^{-1}]$ and t is known as *spectral parameter*.

The following representations, known as *evaluation modules*, form a tensor category of $\hat{\mathfrak{g}}$:

$$V_1(a_1) \otimes V_2(a_2) \otimes \cdots \otimes V_n(a_n),$$

where

- ▶ V_i are representations of \mathfrak{g}
- ▶ a_i are values for t

Let us consider Lie algebra \mathfrak{g} .

The associated *loop algebra* is $\hat{\mathfrak{g}} = \mathfrak{g}[t, t^{-1}]$ and t is known as *spectral parameter*.

The following representations, known as *evaluation modules*, form a tensor category of $\hat{\mathfrak{g}}$:

$$V_1(a_1) \otimes V_2(a_2) \otimes \cdots \otimes V_n(a_n),$$

where

- ▶ V_i are representations of \mathfrak{g}
- ▶ a_i are values for t

Quantum groups:

$$U_{\hbar}(\hat{\mathfrak{g}})$$

are deformations of $U(\hat{\mathfrak{g}})$, with a **nontrivial intertwiner** $R_{V_1, V_2}(a_1/a_2)$:

$$V_1(a_1) \otimes V_2(a_2)$$



$$V_2(a_2) \otimes V_1(a_1)$$

Quantum groups:

$$U_{\hbar}(\hat{\mathfrak{g}})$$

are deformations of $U(\hat{\mathfrak{g}})$, with a **nontrivial intertwiner** $R_{V_1, V_2}(a_1/a_2)$:

$$V_1(a_1) \otimes V_2(a_2)$$



$$V_2(a_2) \otimes V_1(a_1)$$

which is a rational function of a_1, a_2 , satisfying **Yang-Baxter equation**:



The generators of $U_{\hbar}(\hat{\mathfrak{g}})$ emerge as matrix elements of R -matrices:
FRT construction.

Introduction

Quantum Integrable
models

Enumerative geometry
and Bethe ansatz

QQ-systems and
(G, \hbar)-opers

Applications

Physical space:

$$\mathcal{H}_{\text{phys}} = V_1(a_1) \otimes V_2(a_2) \otimes \cdots \otimes V_n(a_n).$$

Auxiliary spaces: $W(u)$.

Quantum monodromy matrix:

$$M(u) = (Z \otimes \text{Id}) \tilde{R}_{W(u), \mathcal{H}_{\text{phys}}} : W(u) \otimes \mathcal{H}_{\text{phys}} \rightarrow W(u) \otimes \mathcal{H}_{\text{phys}}$$

Here \tilde{R} is the R-matrix, composed with permutation operator,
 $Z \in e^{\mathfrak{h}}$ - diagonal.

Physical space:

$$\mathcal{H}_{\text{phys}} = V_1(a_1) \otimes V_2(a_2) \otimes \cdots \otimes V_n(a_n).$$

Auxiliary spaces: $W(u)$.

Quantum monodromy matrix:

$$M(u) = (Z \otimes \text{Id}) \tilde{R}_{W(u), \mathcal{H}_{\text{phys}}} : W(u) \otimes \mathcal{H}_{\text{phys}} \rightarrow W(u) \otimes \mathcal{H}_{\text{phys}}$$

Here \tilde{R} is the R-matrix, composed with permutation operator,
 $Z \in e^{\mathfrak{h}}$ - diagonal.

Physical space:

$$\mathcal{H}_{\text{phys}} = V_1(a_1) \otimes V_2(a_2) \otimes \cdots \otimes V_n(a_n).$$


Auxiliary spaces: $W(u)$.

Quantum monodromy matrix:

$$M(u) = (Z \otimes \text{Id}) \tilde{R}_{W(u), \mathcal{H}_{\text{phys}}} : W(u) \otimes \mathcal{H}_{\text{phys}} \rightarrow W(u) \otimes \mathcal{H}_{\text{phys}}$$

Here \tilde{R} is the R-matrix, composed with permutation operator,
 $Z \in e^{\hbar} \in U_{\hbar}(\hat{\mathfrak{g}})$ - diagonal.

Transfer matrix:



$$T_{W(u)} = \text{Tr}_{W(u)} [M(u)], \quad T_{W(u)} : \mathcal{H}_{\text{phys}} \rightarrow \mathcal{H}_{\text{phys}}$$

Introduction

Quantum Integrable
models

Enumerative geometry
and Bethe ansatz

QQ-systems and
(G, h)-opers

Applications

Integrability:

$$[T_{w'(u')}, T_{w(u)}] = 0$$

follows from Yang-Baxter relation.

Transfer matrices $T_{w(u)}$ generate **Bethe algebra**:

$$T_{w(u)} = \sum_n u^n I_n, \quad [I_n, I_m] = 0.$$

Primary goal: **diagonalize** $\{T_{w(u)}\}$ **simultaneously**.

$\mathfrak{g} = \mathfrak{sl}(2)$: Heisenberg spin chain

Anton M. Zeitlin

Introduction

Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

$$\mathcal{H}_{\text{phys}} = \mathbb{C}^2(a_1) \otimes \mathbb{C}^2(a_2) \otimes \cdots \otimes \mathbb{C}^2(a_n)$$

States : $\uparrow\uparrow\uparrow\uparrow\downarrow\uparrow\uparrow\uparrow\downarrow\uparrow\uparrow\uparrow\uparrow\downarrow\uparrow\uparrow\uparrow\uparrow\downarrow\downarrow\uparrow\uparrow\uparrow$

Here $\mathbb{C}^2(u)$ stands for 2-dimensional representation of $U_{\hbar}(\widehat{\mathfrak{sl}}_2)$.

Algebraic Bethe ansatz as a part of QISM:

$$T_{\mathbb{C}^2(u)} = \text{Tr}_{\mathbb{C}^2(u)} \left[(Z \otimes \text{Id}) \tilde{R}_{\mathbb{C}^2(u), \mathcal{H}_{\text{phys}}} \right] = \text{Tr} \begin{pmatrix} A(u) & B(u) \\ C(u) & D(u) \end{pmatrix} = A(u) + D(u)$$

$$A(u), B(u), C(u), D(u) : \mathcal{H}_{\text{phys}} \rightarrow \mathcal{H}_{\text{phys}}$$

Bethe vectors:

$$|0\rangle = \uparrow\uparrow\uparrow \dots \uparrow\uparrow\uparrow$$

$$\{B(x_1) \dots B(x_k) | 0\rangle; \quad C(x) | 0\rangle = 0\}$$

Commutation relations between A, B, C, D : from Yang-Baxter equation.

$$\mathcal{H}_{\text{phys}} = \mathbb{C}^2(a_1) \otimes \mathbb{C}^2(a_2) \otimes \cdots \otimes \mathbb{C}^2(a_n)$$

States : $\uparrow\uparrow\uparrow\uparrow\downarrow\uparrow\uparrow\uparrow\downarrow\uparrow\uparrow\uparrow\uparrow\downarrow\uparrow\uparrow\uparrow\uparrow\downarrow\downarrow\uparrow\uparrow\uparrow$

Here $\mathbb{C}^2(u)$ stands for 2-dimensional representation of $U_{\hbar}(\widehat{\mathfrak{sl}}_2)$.

Algebraic Bethe ansatz as a part of QISM:

$$T_{\mathbb{C}^2(u)} = \text{Tr}_{\mathbb{C}^2(u)} \left[(Z \otimes \text{Id}) \tilde{R}_{\mathbb{C}^2(u), \mathcal{H}_{\text{phys}}} \right] = \text{Tr} \begin{pmatrix} A(u) & B(u) \\ C(u) & D(u) \end{pmatrix} = A(u) + D(u)$$

$$A(u), B(u), C(u), D(u) : \mathcal{H}_{\text{phys}} \rightarrow \mathcal{H}_{\text{phys}}$$

Bethe vectors:

$$|0\rangle = \uparrow\uparrow\uparrow \dots \uparrow\uparrow\uparrow$$

$$\{B(x_1) \dots B(x_k) | 0\rangle; \quad C(x) | 0\rangle = 0\}$$

Commutation relations between A, B, C, D : from Yang-Baxter equation.

$$\mathcal{H}_{\text{phys}} = \mathbb{C}^2(\mathbf{a}_1) \otimes \mathbb{C}^2(\mathbf{a}_2) \otimes \cdots \otimes \mathbb{C}^2(\mathbf{a}_n)$$

States : 

Here $\mathbb{C}^2(u)$ stands for 2-dimensional representation of $U_h(\widehat{\mathfrak{sl}}_2)$.

Algebraic Bethe ansatz as a part of QISM:

$$T_{\mathbb{C}^2(u)} = \text{Tr}_{\mathbb{C}^2(u)} \left[(Z \otimes \text{Id}) \tilde{R}_{\mathbb{C}^2(u), \mathcal{H}_{\text{phys}}} \right] = \text{Tr} \begin{pmatrix} A(u) & B(u) \\ C(u) & D(u) \end{pmatrix} = A(u) + D(u)$$

$$A(\textcolor{brown}{u}), B(\textcolor{brown}{u}), C(\textcolor{brown}{u}), D(\textcolor{brown}{u}) : \mathcal{H}_{\text{phys}} \rightarrow \mathcal{H}_{\text{phys}}$$

Bethe vectors:

$$|0\rangle = \uparrow\uparrow\uparrow \dots \uparrow\uparrow\uparrow$$

$$\{B(x_1) \dots B(x_k) | 0\rangle; \quad C(x) | 0\rangle = 0\}$$

Commutation relations between A, B, C, D : from Yang-Baxter equation.

The eigenvalues are symmetric functions of **Bethe roots** $\{x_i\}$:

$$\prod_{j=1}^n \frac{x_i - a_j}{\hbar a_j - x_i} = z \hbar^{-n/2} \prod_{\substack{j=1 \\ j \neq i}}^k \frac{x_i \hbar - x_j}{x_i - x_j \hbar}, \quad i = 1, \dots, k,$$

Special element in the Bethe algebra: **Q-operator**.

The eigenvalues $\mathcal{Q}(u)$ of the **Q-operator** are the generating functions for the elementary symmetric functions of Bethe roots:

$$\mathcal{Q}(u) = \prod_{i=1}^k (u - x_i)$$

A real challenge is to describe representation-theoretic meaning of **Q-operator** for general \mathfrak{g} (possibly infinite-dimensional).

The eigenvalues are symmetric functions of **Bethe roots** $\{x_i\}$:

$$\prod_{j=1}^n \frac{x_i - a_j}{\hbar a_j - x_i} = z \hbar^{-n/2} \prod_{\substack{j=1 \\ j \neq i}}^k \frac{x_i \hbar - x_j}{x_i - x_j \hbar}, \quad i = 1, \dots, k,$$

Special element in the Bethe algebra: **Q-operator**.

The eigenvalues $\mathcal{Q}(u)$ of the **Q-operator** are the generating functions for the elementary symmetric functions of Bethe roots:

$$\mathcal{Q}(u) = \prod_{i=1}^k (u - x_i)$$

A real challenge is to describe representation-theoretic meaning of **Q-operator** for general \mathfrak{g} (possibly infinite-dimensional).

Quantum Knizhnik-Zamolodchikov (Frenkel-Reshetikhin) equations:

$$\Psi(a_1, \dots, q a_k, \dots, a_{i_n}, \{z_i\}) = H_k^{(q)} \Psi(a_1, \dots, a_n, \{z_i\}),$$

+

commuting q – difference equations in Z –components (dynamical)

- ▶ Ψ takes values in $\mathcal{H}_{\text{phys}} = V_1(a_1) \otimes V_2(a_2) \otimes \dots \otimes V_n(a_n)$,
- ▶ operators $\{H_i^{(q)}\}$ are expressed in terms of products of R -matrices and twist parameter, e.g.,

$$\Psi(q a_1, \dots, a_n, \{z_i\}) = (Z \otimes 1 \otimes \dots \otimes 1) R_{V_1, V_n} \dots R_{V_1, V_2} \Psi$$

- ▶ $\{H_i^{(1)} = \lim_{q \rightarrow 1} H_i^{(q)}\}$ coincide with transfer matrices of certain kind.

Quantum Knizhnik-Zamolodchikov (Frenkel-Reshetikhin) equations:

$$\Psi(a_1, \dots, qa_k, \dots, a_{i_n}, \{z_i\}) = H_k^{(q)} \Psi(a_1, \dots, a_n, \{z_i\}),$$

+

commuting q – difference equations in Z –components (dynamical)

- ▶ Ψ takes values in $\mathcal{H}_{\text{phys}} = V_1(a_1) \otimes V_2(a_2) \otimes \dots \otimes V_n(a_n)$,
- ▶ operators $\{H_i^{(q)}\}$ are expressed in terms of products of R -matrices and twist parameter, e.g.,

$$\Psi(qa_1, \dots, a_n, \{z_i\}) = (Z \otimes 1 \otimes \dots \otimes 1) R_{V_1, V_n} \dots R_{V_1, V_2} \Psi$$

- ▶ $\{H_i^{(1)} = \lim_{q \rightarrow 1} H_i^{(q)}\}$ coincide with transfer matrices of certain kind.

Types of solutions:

- ▶ jointly analytic in a chamber of $\{a_i\}$:

conformal blocks for $U_{\hbar}(\widehat{\mathfrak{g}})$, i.e. products of Intertwining operators for centrally extended $U_{\hbar}(\widehat{\mathfrak{g}})$, where central charge is related to q .

- ▶ jointly analytic in $\{z_i\}$:

conformal blocks for deformed W -algebra: $W_{q,t}({}^L\mathfrak{g})$.

The relationship between this solutions is an essential part of

Quantum q-Langlands correspondence

M. Aganagic, E. Frenkel, A. Okounkov'17

One can obtain Bethe equations from asymptotic behavior of solutions to qKZ equations in $q \rightarrow 1$ limit.

Types of solutions:

- ▶ jointly analytic in a chamber of $\{a_i\}$:

conformal blocks for $U_{\hbar}(\widehat{\mathfrak{g}})$, i.e. products of Intertwining operators for centrally extended $U_{\hbar}(\widehat{\mathfrak{g}})$, where central charge is related to q .

- ▶ jointly analytic in $\{z_i\}$:

conformal blocks for deformed **W-algebra**: $W_{q,t}({}^L\mathfrak{g})$.

The relationship between this solutions is an essential part of

Quantum q-Langlands correspondence

M. Aganagic, E. Frenkel, A. Okounkov'17

One can obtain Bethe equations from asymptotic behavior of solutions to qKZ equations in $q \rightarrow 1$ limit.

Types of solutions:

- ▶ jointly analytic in a chamber of $\{a_i\}$:

conformal blocks for $U_{\hbar}(\widehat{\mathfrak{g}})$, i.e. products of Intertwining operators for centrally extended $U_{\hbar}(\widehat{\mathfrak{g}})$, where central charge is related to q .

- ▶ jointly analytic in $\{z_i\}$:

conformal blocks for deformed **W-algebra**: $W_{q,t}({}^L\mathfrak{g})$.

The relationship between this solutions is an essential part of

Quantum q-Langlands correspondence

M. Aganagic, E. Frenkel, A. Okounkov'17

One can obtain Bethe equations from asymptotic behavior of solutions to qKZ equations in $q \rightarrow 1$ limit.

Types of solutions:

- ▶ jointly analytic in a chamber of $\{a_i\}$:

conformal blocks for $U_{\hbar}(\widehat{\mathfrak{g}})$, i.e. products of Intertwining operators for centrally extended $U_{\hbar}(\widehat{\mathfrak{g}})$, where central charge is related to q .

- ▶ jointly analytic in $\{z_i\}$:

conformal blocks for deformed **W-algebra**: $W_{q,t}({}^L\mathfrak{g})$.

The relationship between this solutions is an essential part of

Quantum q-Langlands correspondence

M. Aganagic, E. Frenkel, A. Okounkov'17

One can obtain Bethe equations from asymptotic behavior of solutions to qKZ equations in $q \rightarrow 1$ limit.

Types of solutions:

- ▶ jointly analytic in a chamber of $\{a_i\}$:

conformal blocks for $U_{\hbar}(\widehat{\mathfrak{g}})$, i.e. products of Intertwining operators for centrally extended $U_{\hbar}(\widehat{\mathfrak{g}})$, where central charge is related to q .

- ▶ jointly analytic in $\{z_i\}$:

conformal blocks for deformed **W-algebra**: $W_{q,t}({}^L\mathfrak{g})$.

The relationship between this solutions is an essential part of

Quantum q-Langlands correspondence

M. Aganagic, E. Frenkel, A. Okounkov'17

One can obtain Bethe equations from asymptotic behavior of solutions to qKZ equations in $q \rightarrow 1$ limit.

\mathcal{M} : affine space constructed via quiver (oriented graph) which is the representation of

$$G = GL(v_1) \times GL(v_2) \cdots \times GL(v_{\text{rank}(\mathfrak{g})}),$$

We build \mathcal{M} as a direct sum of:

- ▶ $\oplus_i \text{Hom}(V_i, W_i)$, where $\dim(V_i) = v_i$, W_i is known as **framing**
- ▶ $\oplus_{i \rightarrow j} \text{Hom}(V_i, V_j)$

$T^*\mathcal{M}$: phase space with Poisson bracket.

Nakajima quiver variety is a “clever” quotient, called **algebraic symplectic reduction**:

$$N = T^*\mathcal{M} // G$$

Nakajima, Varagnolo-Vasserot, Maulik-Okounkov: Localized equivariant cohomology/K-theory of N has the structure of weight subspace for representations of $Y_{\hbar}(\mathfrak{g})/U_{\hbar}(\mathfrak{g})$. Weight is determined by a collection $v_1, \dots, v_{\text{rank}(\mathfrak{g})}$

[Introduction](#)[Quantum Integrable models](#)[Enumerative geometry and Bethe ansatz](#)[QQ-systems and \$\(G, \hbar\)\$ -opers](#)[Applications](#)

\mathcal{M} : affine space constructed via quiver (oriented graph) which is the representation of

$$G = GL(v_1) \times GL(v_2) \cdots \times GL(v_{\text{rank}(\mathfrak{g})}),$$

We build \mathcal{M} as a direct sum of:

- ▶ $\oplus_i \text{Hom}(V_i, W_i)$, where $\dim(V_i) = v_i$, W_i is known as **framing**
- ▶ $\oplus_{i \rightarrow j} \text{Hom}(V_i, V_j)$

$T^*\mathcal{M}$: phase space with Poisson bracket.

Nakajima quiver variety is a “clever” quotient, called **algebraic symplectic reduction**:

$$N = T^*\mathcal{M} // G$$

Nakajima, Varagnolo-Vasserot, Maulik-Okounkov: Localized equivariant cohomology/K-theory of N has the structure of weight subspace for representations of $Y_{\hbar}(\mathfrak{g})/U_{\hbar}(\mathfrak{g})$. Weight is determined by a collection $v_1, \dots, v_{\text{rank}(\mathfrak{g})}$

[Introduction](#)[Quantum Integrable models](#)[Enumerative geometry and Bethe ansatz](#)[QQ-systems and \$\(G, \hbar\)\$ -opers](#)[Applications](#)

\mathcal{M} : affine space constructed via quiver (oriented graph) which is the representation of

$$G = GL(v_1) \times GL(v_2) \cdots \times GL(v_{\text{rank}(\mathfrak{g})}),$$

We build \mathcal{M} as a direct sum of:

- ▶ $\oplus_i \text{Hom}(V_i, W_i)$, where $\dim(V_i) = v_i$, W_i is known as **framing**
- ▶ $\oplus_{i \rightarrow j} \text{Hom}(V_i, V_j)$

$T^*\mathcal{M}$: phase space with Poisson bracket.

Nakajima quiver variety is a “clever” quotient, called **algebraic symplectic reduction**:

$$N = T^*\mathcal{M} // G$$

Nakajima, Varagnolo-Vasserot, Maulik-Okounkov: Localized equivariant cohomology/K-theory of N has the structure of weight subspace for representations of $Y_{\hbar}(\mathfrak{g})/U_{\hbar}(\mathfrak{g})$. Weight is determined by a collection $v_1, \dots, v_{\text{rank}(\mathfrak{g})}$

[Introduction](#)[Quantum Integrable models](#)[Enumerative geometry and Bethe ansatz](#)[QQ-systems and \$\(G, \hbar\)\$ -opers](#)[Applications](#)

\mathcal{M} : affine space constructed via quiver (oriented graph) which is the representation of

$$G = GL(v_1) \times GL(v_2) \cdots \times GL(v_{\text{rank}(\mathfrak{g})}),$$

We build \mathcal{M} as a direct sum of:

- ▶ $\oplus_i \text{Hom}(V_i, W_i)$, where $\dim(V_i) = v_i$, W_i is known as **framing**
- ▶ $\oplus_{i \rightarrow j} \text{Hom}(V_i, V_j)$

$T^*\mathcal{M}$: phase space with Poisson bracket.

Nakajima quiver variety is a “clever” quotient, called **algebraic symplectic reduction**:

$$N = T^*\mathcal{M} // G$$

Nakajima, Varagnolo-Vasserot, Maulik-Okounkov: Localized equivariant cohomology/K-theory of N has the structure of weight subspace for representations of $Y_{\hbar}(\mathfrak{g})/U_{\hbar}(\mathfrak{g})$. Weight is determined by a collection $v_1, \dots, v_{\text{rank}(\mathfrak{g})}$

[Introduction](#)[Quantum Integrable models](#)[Enumerative geometry and Bethe ansatz](#)[QQ-systems and \$\(G, \hbar\)\$ -opers](#)[Applications](#)

Example: $N_{k,n} = T^*Gr(k, n)$

Anton M. Zeitlin

Introduction

Quantum Integrable
models

Enumerative geometry
and Bethe ansatz

QQ-systems and
(G, \hbar)-opers

Applications

Let $\dim(V) = k$, $\dim(W) = n$,

$$\mathcal{M} = \operatorname{Hom}(V, W),$$

$$T^*\mathcal{M} = \operatorname{Hom}(V, W) \oplus \operatorname{Hom}(W, V)$$

The **moment map**:

$$\mu : T^*\mathcal{M} \rightarrow \operatorname{End}(V); \quad (A, B) \mapsto BA$$

generates the $GL(V)$ -action.

Then

$$T^*Gr(k, n) = N_{k,n} = T^*\mathcal{M} // GL(V) = \mu^{-1}(0) // GL(V) = \mu^{-1}(0)_{ss} / GL(V).$$

The semi-stability condition: $\operatorname{Hom}(V, W)$ is injective.

Example: $N_{k,n} = T^*Gr(k, n)$

Anton M. Zeitlin

Introduction

Quantum Integrable
models

Enumerative geometry
and Bethe ansatz

QQ-systems and
(G, \hbar)-opers

Applications

Let $\dim(V) = k$, $\dim(W) = n$,

$$\mathcal{M} = \operatorname{Hom}(V, W),$$

$$T^*\mathcal{M} = \operatorname{Hom}(V, W) \oplus \operatorname{Hom}(W, V)$$

The **moment map**:

$$\mu : T^*\mathcal{M} \rightarrow \operatorname{End}(V); \quad (A, B) \mapsto BA$$

generates the $GL(V)$ -action.

Then

$$T^*Gr(k, n) = N_{k,n} = T^*\mathcal{M} // GL(V) = \mu^{-1}(0) // GL(V) = \mu^{-1}(0)_{ss} / GL(V).$$

The semi-stability condition: $\operatorname{Hom}(V, W)$ is injective.

Example: $N_{k,n} = T^*Gr(k, n)$

Anton M. Zeitlin

Introduction

Quantum Integrable
models

Enumerative geometry
and Bethe ansatz

QQ-systems and
(G, \hbar)-opers

Applications

Let $\dim(V) = k$, $\dim(W) = n$,

$$\mathcal{M} = \operatorname{Hom}(V, W),$$

$$T^*\mathcal{M} = \operatorname{Hom}(V, W) \oplus \operatorname{Hom}(W, V)$$

The **moment map**:

$$\mu : T^*\mathcal{M} \rightarrow \operatorname{End}(V); \quad (A, B) \mapsto BA$$

generates the $GL(V)$ -action.

Then

$$T^*Gr(k, n) = N_{k,n} = T^*\mathcal{M} // GL(V) = \mu^{-1}(0) // GL(V) = \mu^{-1}(0)_{ss} / GL(V).$$

The semi-stability condition: $\operatorname{Hom}(V, W)$ is injective.

Torus action:

$$A = \mathbb{C}_{a_1}^\times \times \cdots \times \mathbb{C}_{a_n}^\times \curvearrowright W,$$

Full torus : $T = A \times \mathbb{C}_\hbar^\times$, where \mathbb{C}_\hbar^\times scales cotangent directions

Tautological bundles:

$$\mathcal{V} = T^*\mathcal{M} \times V // GL(V), \quad \mathcal{W} = T^*\mathcal{M} \times W // GL(V)$$

For any $\tau \in S(x_1^{\pm 1}, x_2^{\pm 1}, \dots, x_k^{\pm 1})$ we have:

$$\tau(\mathcal{V}) = T^*\mathcal{M} \times \tau(V) // GL(V)$$

Example: $\tau(V) = V^{\otimes 2} - \Lambda^3 V^*$ corresponds to

$$\tau(x_1, \dots, x_k) = (x_1 + \cdots + x_k)^2 - \sum_{1 \leq i_1 \leq i_2 \leq i_3 \leq k} x_{i_1}^{-1} x_{i_2}^{-1} x_{i_3}^{-1}.$$

Equivariant K-theory is generated by tautological bundles under tensor product.

Introduction

Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

Torus action:

$$A = \mathbb{C}_{a_1}^\times \times \cdots \times \mathbb{C}_{a_n}^\times \curvearrowright W,$$

Full torus : $T = A \times \mathbb{C}_\hbar^\times$, where \mathbb{C}_\hbar^\times scales cotangent directions

Tautological bundles:

$$\mathcal{V} = T^*\mathcal{M} \times V // GL(V), \quad \mathcal{W} = T^*\mathcal{M} \times W // GL(V)$$

For any $\tau \in S(x_1^{\pm 1}, x_2^{\pm 1}, \dots, x_k^{\pm 1})$ we have:

$$\tau(\mathcal{V}) = T^*\mathcal{M} \times \tau(V) // GL(V)$$

Example: $\tau(V) = V^{\otimes 2} - \Lambda^3 V^*$ corresponds to

$$\tau(x_1, \dots, x_k) = (x_1 + \cdots + x_k)^2 - \sum_{1 \leq i_1 \leq i_2 \leq i_3 \leq k} x_{i_1}^{-1} x_{i_2}^{-1} x_{i_3}^{-1}.$$

Equivariant K-theory is generated by tautological bundles under tensor product.

Introduction

Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

Torus action:

$$A = \mathbb{C}_{a_1}^\times \times \cdots \times \mathbb{C}_{a_n}^\times \curvearrowright W,$$

Full torus : $T = A \times \mathbb{C}_\hbar^\times$, where \mathbb{C}_\hbar^\times scales cotangent directions

Tautological bundles:

$$\mathcal{V} = T^*\mathcal{M} \times V // GL(V), \quad \mathcal{W} = T^*\mathcal{M} \times W // GL(V)$$

For any $\tau \in S(x_1^{\pm 1}, x_2^{\pm 1}, \dots, x_k^{\pm 1})$ we have:

$$\tau(\mathcal{V}) = T^*\mathcal{M} \times \tau(V) // GL(V)$$

Example: $\tau(V) = V^{\otimes 2} - \Lambda^3 V^*$ corresponds to

$$\tau(x_1, \dots, x_k) = (x_1 + \cdots + x_k)^2 - \sum_{1 \leq i_1 \leq i_2 \leq i_3 \leq k} x_{i_1}^{-1} x_{i_2}^{-1} x_{i_3}^{-1}.$$

Equivariant K-theory is generated by tautological bundles under tensor product.

Introduction

Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

Torus action:

$$A = \mathbb{C}_{a_1}^\times \times \cdots \times \mathbb{C}_{a_n}^\times \curvearrowright W,$$

Full torus : $T = A \times \mathbb{C}_h^\times$, where \mathbb{C}_h^\times scales cotangent directions

Tautological bundles:

$$\mathcal{V} = T^*\mathcal{M} \times V // GL(V), \quad \mathcal{W} = T^*\mathcal{M} \times W // GL(V)$$

For any $\tau \in S(x_1^{\pm 1}, x_2^{\pm 1}, \dots, x_k^{\pm 1})$ we have:

$$\tau(\mathcal{V}) = T^*\mathcal{M} \times \tau(V) // GL(V)$$

Example: $\tau(V) = V^{\otimes 2} - \Lambda^3 V^*$ corresponds to

$$\tau(x_1, \dots, x_k) = (x_1 + \cdots + x_k)^2 - \sum_{1 \leq i_1 \leq i_2 \leq i_3 \leq k} x_{i_1}^{-1} x_{i_2}^{-1} x_{i_3}^{-1}.$$

Equivariant K-theory is generated by tautological bundles under tensor product.

Introduction

Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

Torus action:

$$A = \mathbb{C}_{a_1}^\times \times \cdots \times \mathbb{C}_{a_n}^\times \curvearrowright W,$$

Full torus : $\mathcal{T} = A \times \mathbb{C}_\hbar^\times$, where \mathbb{C}_\hbar^\times scales cotangent directions

Fixed points: $\mathbf{p} = \{s_1, \dots, s_k\} \in \{a_1, \dots, a_n\}$

Let $N(n) = \sqcup_k N_{k,n} = \sqcup_k T^* Gr(k, n)$.

Localized K-theory: $K_T(N(n))_{loc}$ as a $\mathbb{Q}(a_1, \dots, a_n, \hbar)$ -module is identified with:

$$\mathcal{H}_{\text{phys}} = \mathbb{C}^2(a_1) \otimes \mathbb{C}^2(a_2) \otimes \cdots \otimes \mathbb{C}^2(a_n)$$

generated by $\mathcal{O}_{\mathbf{p}}$.

“Classical” Bethe equations: The eigenvalues of the operators of multiplication by τ are $\tau(x_1, \dots, x_k)$ evaluated at the solutions of the following equations:

$$\prod_{j=1}^n (x_i - a_j) = 0, \quad i = 1, \dots, k, \text{ with } x_i \neq x_j$$

Introduction

Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

Torus action:

$$A = \mathbb{C}_{a_1}^\times \times \cdots \times \mathbb{C}_{a_n}^\times \curvearrowright W,$$

Full torus : $\mathcal{T} = \mathcal{A} \times \mathbb{C}_h^\times$, where \mathbb{C}_h^\times scales cotangent directions

Fixed points: $\mathbf{p} = \{s_1, \dots, s_k\} \in \{a_1, \dots, a_n\}$

Let $N(n) = \sqcup_k N_{k,n} = \sqcup_k T^* Gr(k, n)$.

Localized K-theory: $K_T(N(n))_{loc}$ as a $\mathbb{Q}(a_1, \dots, a_n, \hbar)$ -module is identified with:

$$\mathcal{H}_{\text{phys}} = \mathbb{C}^2(\mathbf{a}_1) \otimes \mathbb{C}^2(\mathbf{a}_2) \otimes \cdots \otimes \mathbb{C}^2(\mathbf{a}_n)$$

generated by $\mathcal{O}_{\mathbf{p}}$.

Torus action:

$$A = \mathbb{C}_{a_1}^\times \times \cdots \times \mathbb{C}_{a_n}^\times \curvearrowright W,$$

Full torus : $T = A \times \mathbb{C}_{\hbar}^\times$, where $\mathbb{C}_{\hbar}^\times$ scales cotangent directions

Fixed points: $\mathbf{p} = \{s_1, \dots, s_k\} \in \{a_1, \dots, a_n\}$

Let $N(n) = \sqcup_k N_{k,n} = \sqcup_k T^*Gr(k, n)$.

Localized K-theory: $K_T(N(n))_{loc}$ as a $\mathbb{Q}(a_1, \dots, a_n, \hbar)$ -module is identified with:

$$\mathcal{H}_{\text{phys}} = \mathbb{C}^2(a_1) \otimes \mathbb{C}^2(a_2) \otimes \cdots \otimes \mathbb{C}^2(a_n)$$

generated by $\mathcal{O}_{\mathbf{p}}$.

“Classical” Bethe equations: The eigenvalues of the operators of multiplication by τ are $\tau(x_1, \dots, x_k)$ evaluated at the solutions of the following equations:

$$\prod_{j=1}^n (x_i - a_j) = 0, \quad i = 1, \dots, k, \text{ with } x_i \neq x_j$$

Introduction

Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

Quasimap $f: \mathcal{C} = \mathbb{P}^1 \dashrightarrow N_{k,n}$ is the following collection of data:

- ▶ vector bundle \mathcal{V} on \mathbb{P}^1 of rank k .
- ▶ section $f \in H^0(\mathbb{P}^1, \mathcal{M} \oplus \mathcal{M}^* \otimes \hbar)$, satisfying the condition $\mu = 0$, where $\mathcal{M} = \text{Hom}(\mathcal{V}, \mathcal{W})$, so that \mathcal{W} is a trivial bundle of rank n .

$$\text{ev}_p(f) = f(p) \in [\mu^{-1}(0)/GL(V)] \supset N_{k,n}$$

Quasimap is *stable* if $f(p) \in N_{k,n}$ for all but finitely many points, known as *singularities* of quasimap.

For the moduli space of stable quasimaps

$$QM(N_{k,n})$$

only \mathcal{V} and f vary, while \mathcal{C} and \mathcal{W} remain the same.

$$\deg(f) := \deg(\mathcal{V}), \quad QM(N_{k,n}) = \sqcup_{d \geq 0} QM^d(N_{k,n}).$$

Quasimap $f: \mathcal{C} = \mathbb{P}^1 \dashrightarrow N_{k,n}$ is the following collection of data:

- ▶ vector bundle \mathcal{V} on \mathbb{P}^1 of rank k .
- ▶ section $f \in H^0(\mathbb{P}^1, \mathcal{M} \oplus \mathcal{M}^* \otimes \hbar)$, satisfying the condition $\mu = 0$, where $\mathcal{M} = \text{Hom}(\mathcal{V}, \mathcal{W})$, so that \mathcal{W} is a trivial bundle of rank n .

$$\text{ev}_p(f) = f(p) \in [\mu^{-1}(0)/GL(V)] \supset N_{k,n}$$

Quasimap is *stable* if $f(p) \in N_{k,n}$ for all but finitely many points, known as *singularities* of quasimap.

For the moduli space of stable quasimaps

$$QM(N_{k,n})$$

only \mathcal{V} and f vary, while \mathcal{C} and \mathcal{W} remain the same.

$$\deg(f) := \deg(\mathcal{V}), \quad QM(N_{k,n}) = \sqcup_{d \geq 0} QM^d(N_{k,n}).$$

Quasimap $f: \mathcal{C} = \mathbb{P}^1 \dashrightarrow N_{k,n}$ is the following collection of data:

- ▶ vector bundle \mathcal{V} on \mathbb{P}^1 of rank k .
- ▶ section $f \in H^0(\mathbb{P}^1, \mathcal{M} \oplus \mathcal{M}^* \otimes \hbar)$, satisfying the condition $\mu = 0$, where $\mathcal{M} = \text{Hom}(\mathcal{V}, \mathcal{W})$, so that \mathcal{W} is a trivial bundle of rank n .

$$\text{ev}_p(f) = f(p) \in [\mu^{-1}(0)/GL(V)] \supset N_{k,n}$$

Quasimap is *stable* if $f(p) \in N_{k,n}$ for all but finitely many points, known as *singularities* of quasimap.

For the moduli space of stable quasimaps

$$QM(N_{k,n})$$

only \mathcal{V} and f vary, while \mathcal{C} and \mathcal{W} remain the same.

$$\deg(f) := \deg(\mathcal{V}), \quad QM(N_{k,n}) = \sqcup_{d \geq 0} QM^d(N_{k,n}).$$


Introduction

Quantum Integrable models


Enumerative geometry and Bethe ansatz

QQ-systems and
(G, \hbar)-opers

Applications




In 2009 Nekrasov and Shatashvili looked at 3d SUSY gauge theories on $\mathbb{C} \times S^1$:



Gauge field theory with gauge group $G = \times_{i=1}^{rank g} U(v_i)$ defined by a certain action functional $S(\phi_{\{\alpha\}}, A_{\{i\}})$.

- ▶ $A_{\{i\}}$: $U(v_i)$ -connections (gauge fields)
- ▶ $\phi_{\{\alpha\}}$: sections of associated vector $U(v_i)$ -bundles, corresponding to the quiver data (matter fields)

In 2009 Nekrasov and Shatashvili looked at 3d SUSY gauge theories on $\mathbb{C} \times S^1$:




Gauge field theory with gauge group $G = \times_{i=1}^{rank g} U(v_i)$ defined by a certain action functional $S(\phi_{\{\alpha\}}, A_{\{i\}})$.

- ▶ $A_{\{i\}}$: $U(v_i)$ -connections (gauge fields)
- ▶ $\phi_{\{\alpha\}}$: sections of associated vector $U(v_i)$ -bundles, corresponding to the quiver data (matter fields)

Physicists compute **path integrals**:

$$\langle F \rangle = \int [d\phi_{\{\alpha\}}][dA_{\{i\}}] e^{-S(\phi_{\{\alpha\}}, A_{\{i\}})} F(\phi_{\{\alpha\}}, A_{\{i\}})$$

In 2009 Nekrasov and Shatashvili looked at 3d SUSY gauge theories on $\mathbb{C} \times S^1$:



Gauge field theory with gauge group $G = \times_{i=1}^{rank g} U(v_i)$ defined by a certain action functional $S(\phi_{\{\alpha\}}, A_{\{i\}})$.

- ▶ $A_{\{i\}}$: $U(v_i)$ -connections (gauge fields)
- ▶ $\phi_{\{\alpha\}}$: sections of associated vector $U(v_i)$ -bundles, corresponding to the quiver data (matter fields)

Physicists compute **path integrals**:

$$\langle F \rangle = \int [d\phi_{\{\alpha\}}][dA_{\{i\}}] e^{-S(\phi_{\{\alpha\}}, A_{\{i\}})} F(\phi_{\{\alpha\}}, A_{\{i\}})$$

Minima of S : Moduli of Higgs vacua \longleftrightarrow Nakajima quiver variety:

$$T^*\mathcal{M} \mathbin{////} G = \mu^{-1}(0) \mathbin{//} G = N$$


where $\mu = 0$ is a momentum map (low energy configuration) condition.

$\langle F \rangle$ corresponds to weighted K-theoretic counts of **quasimaps**:

Euler characteristics of equivariant pushforwards of sheaves of quasimap moduli space $QM(N_{k,n})$.

The weight (Kähler) parameter is $Z^{\deg(f)}$, which is exactly twist parameter Z we encountered before.

“Line operators”—traces of holonomy of gauge fields generate **quantum K-theory ring**, so that the structure constants of algebra are given by:




$\langle F \rangle$ corresponds to weighted K-theoretic counts of **quasimaps**:

Euler characteristics of equivariant pushforwards of sheaves of quasimap moduli space $QM(N_{k,n})$.

The weight (Kähler) parameter is $z^{\deg(f)}$, which is exactly twist parameter z we encountered before.

“Line operators”—traces of holonomy of gauge fields generate **quantum K-theory ring**, so that the structure constants of algebra are given by:



Conjecture of Nekrasov and Shatashvili '09 (through 3D gauge theory):

Quantum equivariant K – theory ring of quiver variety =

Bethe algebra of related spin chain system

Introduction

Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and
(G, \hbar)-opers

Applications

Quasimaps to quiver varieties:

A diagram consisting of a black circle. Inside the circle is a red curved arrow pointing from the bottom-left towards the top-right, with a red 'q' written below it. To the right of the circle is a dashed green arrow pointing to the right, towards the letter 'N'.

Deformation of the product: $A \circ B = A \otimes B + \sum_{d=1}^{\infty} A \otimes_d B z^d$.

Quantum tautological classes – deformations of $\tau = T^*\mathcal{M} \times \tau(V)$:

$$\hat{\tau}(z) = \tau + \sum_{d=1}^{\infty} \tau_d z^d \in K_T(N(n))[[z]]$$

Theorem.

- ▶ The eigenvalues of operators of quantum multiplication by $\hat{\tau}(z)$ are given by the values of the corresponding Laurent polynomials $\tau(x_1, \dots, x_k)$ evaluated at the solutions of Bethe equations:

$$\prod_{j=1}^n \frac{x_i - a_j}{\hbar a_j - x_i} = z \hbar^{-n/2} \prod_{\substack{j=1 \\ i \neq j}}^k \frac{x_i \hbar - x_j}{x_i - x_j \hbar}, \quad i = 1, \dots, k,$$

- Baxter Q-operator: $Q(u) = \sum_{i=1}^k (-1)^i u^{k-i} [\Lambda^i V](z) \circledast$

P. Pushkar, A. Smirnov, A.Z.,

Baxter Q-operator from quantum K-theory, Adv. Math. '20

- Generalization to type A_n , also connection to classical many body-systems for T^*FI .

P. Koroteev, P. Pushkar, A. Smirnov, A.Z.,
Quantum K-theory of Quiver Varieties and Many-Body Systems,
Selecta Math. '21

- Connection to quantum many-body systems (as in [Givental-Lee](#)) in the context of partial flags through the works of Cherednik and Matsuo.

P. Koroteev, A.Z.,
qKZ/tRS Duality via Quantum K-Theoretic Counts,
Math. Res. Lett. '21

- 3D mirror symmetry for instanton spaces and their cyclic quiver generalizations

P. Koroteev, A. Z.,
3d Mirror Symmetry for Instanton Moduli Spaces,
Comm. Math. Phys. '23

- Generalization to type A_n , also connection to classical many body-systems for T^*FI .

P. Koroteev, P. Pushkar, A. Smirnov, A.Z.,
Quantum K-theory of Quiver Varieties and Many-Body Systems,
Selecta Math. '21

- Connection to quantum many-body systems (as in [Givental-Lee](#)) in the context of partial flags through the works of Cherednik and Matsuo.

P. Koroteev, A.Z.,
qKZ/tRS Duality via Quantum K-Theoretic Counts,
Math. Res. Lett. '21

- 3D mirror symmetry for instanton spaces and their cyclic quiver generalizations

P. Koroteev, A. Z.,
3d Mirror Symmetry for Instanton Moduli Spaces,
Comm. Math. Phys. '23

- Generalization to type A_n , also connection to classical many body-systems for T^*FI .

P. Koroteev, P. Pushkar, A. Smirnov, A.Z.,
Quantum K-theory of Quiver Varieties and Many-Body Systems,
Selecta Math. '21

- Connection to quantum many-body systems (as in [Givental-Lee](#)) in the context of partial flags through the works of Cherednik and Matsuo.

P. Koroteev, A.Z.,
qKZ/tRS Duality via Quantum K-Theoretic Counts,
Math. Res. Lett. '21

- 3D mirror symmetry for instanton spaces and their cyclic quiver generalizations

P. Koroteev, A. Z.,
3d Mirror Symmetry for Instanton Moduli Spaces,
Comm. Math. Phys. '23

Short exact sequence of bundles:

$$0 \rightarrow V \rightarrow W \rightarrow \hbar \otimes V^\vee \rightarrow 0$$

$$Q(u) = \sum_{i=1}^k (-1)^i u^{k-i} [\wedge^i V](z) \circledast$$

$$\tilde{Q}(u) = \sum_{i=1}^k (-1)^i u^{k-i} [\wedge^i V^\vee](z) \circledast$$

Eigenvalues of these two operators are related via **QQ-system**:

$$\tilde{Q}(\hbar u) Q(u) - z Q(\hbar u) \tilde{Q}(u) = \prod_i (u - a_i)$$

Equivalent to Bethe ansatz equations.

$$M_{\hbar} : \mathbb{P}^1 \rightarrow \mathbb{P}^1, \text{ such that } u \rightarrow \hbar u.$$

Let \mathcal{F}_G be a G -bundle over \mathbb{P}^1 :

- ▶ G - simple simply connected Lie group associated to Lie algebra \mathfrak{g} .
- ▶ \mathcal{F}_G^{\hbar} is a pushforward w.r.t. M_{\hbar} .

(G, \hbar) -connection: a meromorphic section of $\text{Hom}_{\mathcal{O}_{\mathbb{P}^1}}(\mathcal{F}_G, \mathcal{F}_G^{\hbar})$.

Locally:

\hbar -gauge transformations of (G, \hbar) -connection:

$$A(u) \rightarrow g(\hbar u) A(u) g^{-1}(u),$$

where $A(u) \in G(u) = G(\mathbb{C}(u))$.

Compare it with standard gauge transformations:

$$\partial_u + A(u) \rightarrow g(u)(\partial_u + A(u))g^{-1}(u),$$

where $A(u) \in \mathfrak{g}(u)$.

$$M_{\hbar} : \mathbb{P}^1 \rightarrow \mathbb{P}^1, \text{ such that } u \rightarrow \hbar u.$$

Let \mathcal{F}_G be a G -bundle over \mathbb{P}^1 :

- ▶ G - simple simply connected Lie group associated to Lie algebra \mathfrak{g} .
- ▶ \mathcal{F}_G^{\hbar} is a pushforward w.r.t. M_{\hbar} .

(G, \hbar) -connection: a meromorphic section of $\text{Hom}_{\mathcal{O}_{\mathbb{P}^1}}(\mathcal{F}_G, \mathcal{F}_G^{\hbar})$.

Locally:

\hbar -gauge transformations of (G, \hbar) -connection:

$$A(u) \rightarrow g(\hbar u)A(u)g^{-1}(u),$$

where $A(u) \in G(u) = G(\mathbb{C}(u))$.

Compare it with standard gauge transformations:

$$\partial_u + A(u) \rightarrow g(u)(\partial_u + A(u))g^{-1}(u),$$

where $A(u) \in \mathfrak{g}(u)$.

$$M_{\hbar} : \mathbb{P}^1 \rightarrow \mathbb{P}^1, \text{ such that } u \rightarrow \hbar u.$$

Let \mathcal{F}_G be a G -bundle over \mathbb{P}^1 :

- ▶ G - simple simply connected Lie group associated to Lie algebra \mathfrak{g} .
- ▶ \mathcal{F}_G^{\hbar} is a pushforward w.r.t. M_{\hbar} .

(G, \hbar) -connection: a meromorphic section of $\text{Hom}_{\mathcal{O}_{\mathbb{P}^1}}(\mathcal{F}_G, \mathcal{F}_G^{\hbar})$.

Locally:

\hbar -gauge transformations of (G, \hbar) -connection:

$$A(u) \rightarrow g(\hbar u)A(u)g^{-1}(u),$$

where $A(u) \in G(u) = G(\mathbb{C}(u))$.

Compare it with standard gauge transformations:

$$\partial_u + A(u) \rightarrow g(u)(\partial_u + A(u))g^{-1}(u),$$

where $A(u) \in \mathfrak{g}(u)$.

$$M_{\hbar} : \mathbb{P}^1 \rightarrow \mathbb{P}^1, \text{ such that } u \rightarrow \hbar u.$$

Let \mathcal{F}_G be a G -bundle over \mathbb{P}^1 :

- ▶ G - simple simply connected Lie group associated to Lie algebra \mathfrak{g} .
- ▶ \mathcal{F}_G^{\hbar} is a pushforward w.r.t. M_{\hbar} .

(G, \hbar) -connection: a meromorphic section of $\text{Hom}_{\mathcal{O}_{\mathbb{P}^1}}(\mathcal{F}_G, \mathcal{F}_G^{\hbar})$.

Locally:

\hbar -gauge transformations of (G, \hbar) -connection:

$$A(u) \rightarrow g(\hbar u)A(u)g^{-1}(u),$$

where $A(u) \in G(u) = G(\mathbb{C}(u))$.

Compare it with standard gauge transformations:

$$\partial_u + A(u) \rightarrow g(u)(\partial_u + A(u))g^{-1}(u),$$

where $A(u) \in \mathfrak{g}(u)$.

Miura (G, \hbar) -oper is a quadruple : $(\mathcal{F}_G, A, \mathcal{F}_{B_+}, \mathcal{F}_{B_-})$, such that:

- ▶ A is (G, \hbar) - connection
- ▶ **Oper condition**: \mathcal{F}_{B_+} : A lies in the Coxeter cell B_+cB_+
- ▶ **Miura condition**: \mathcal{F}_{B_-} : preserved by A

(G, \hbar) -oper is **Z -twisted** if it is gauge equivalent to $Z \in H$, namely

$$A(u) = v(\hbar u) Z v^{-1}(u), \text{ where } Z = \prod_i \zeta_i^{\check{\alpha}_i} \in H, \quad v(u) \in G(u).$$

Example: $SL(r+1)$

$$A \in \begin{bmatrix} * & 0 & 0 & \cdots & 0 \\ * & * & 0 & \cdots & 0 \\ 0 & * & * & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & * & * \end{bmatrix} \quad Z - \text{diagonal}$$

Regular singularities: $\{\Lambda_i(u)\}_{i=1}^r$ -polynomials on subdiagonal.

Introduction

- ## Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

◀ ◻ ▶ ◀ ◼ ▶ ◀ ≡ ▶ ◀ ≡ ▶ ≡

Miura (G, \hbar) -oper is a quadruple : $(\mathcal{F}_G, A, \mathcal{F}_{B_+}, \mathcal{F}_{B_-})$, such that:

- ▶ A is (G, \hbar) - connection
- ▶ **Oper condition:** \mathcal{F}_{B_+} : A lies in the Coxeter cell $B_+ c B_+$
- ▶ **Miura condition:** \mathcal{F}_B : preserved by A

Example: $SL(r+1)$

Miura (G, \hbar) -oper is a quadruple : $(\mathcal{F}_G, A, \mathcal{F}_{B_+}, \mathcal{F}_{B_-})$, such that:

- ▶ A is (G, \hbar) - connection
- ▶ **Oper condition:** \mathcal{F}_{B_+} : A lies in the Coxeter cell B_+cB_+
- ▶ **Miura condition:** \mathcal{F}_B : preserved by A

(G, \hbar) -oper is **Z-twisted** if it is gauge equivalent to $Z \in H$, namely

$$A(u) = v(\hbar u)Zv^{-1}(u), \text{ where } Z = \prod_i \zeta_i^{\check{\alpha}_i} \in H, \ v(u) \in G(u).$$

Example: $SL(r + 1)$

Miura (G, \hbar) -oper is a quadruple $(\mathcal{F}_G, A, \mathcal{F}_{B_+}, \mathcal{F}_{B_-})$, such that:

- ▶ A is (G, \hbar) - connection
- ▶ **Oper condition**: \mathcal{F}_{B_+} : A lies in the Coxeter cell B_+cB_+
- ▶ **Miura condition**: \mathcal{F}_{B_-} : preserved by A

(G, \hbar) -oper is **Z -twisted** if it is gauge equivalent to $Z \in H$, namely

$$A(u) = v(\hbar u) Z v^{-1}(u), \text{ where } Z = \prod_i \zeta_i^{\check{\alpha}_i} \in H, \quad v(u) \in G(u).$$

Example: $SL(r+1)$

$$A \in \begin{bmatrix} * & 0 & 0 & \cdots & 0 \\ * & * & 0 & \cdots & 0 \\ 0 & * & * & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & * & * \end{bmatrix} \quad Z - \text{diagonal}$$

Regular singularities: $\{\Lambda_i(u)\}_{i=1}^r$ -polynomials on subdiagonal.

Miura (G, \hbar) -oper is a quadruple : $(\mathcal{F}_G, A, \mathcal{F}_{B_+}, \mathcal{F}_{B_-})$, such that:

- ▶ A is (G, \hbar) - connection
- ▶ **Oper condition:** \mathcal{F}_{B_+} : A lies in the Coxeter cell $B_+ c B_+$
- ▶ **Miura condition:** \mathcal{F}_B : preserved by A

(G, \hbar) -oper is **Z -twisted** if it is gauge equivalent to $Z \in H$, namely

$$A(u) = v(\hbar u)Zv^{-1}(u), \text{ where } Z = \prod_i \zeta_i^{\check{\alpha}_i} \in H, \ v(u) \in G(u).$$

Example: $SL(r + 1)$

$$A \in \begin{bmatrix} * & 0 & 0 & \cdots & 0 \\ * & * & 0 & \cdots & 0 \\ 0 & * & * & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & * & * \end{bmatrix} \quad \text{Z-diagonal}$$

Regular singularities: $\{\Lambda_i(u)\}_{i=1}^r$ -polynomials on subdiagonal.

Theorem.

There is a one-to-one correspondence between the set of nondegenerate Z -twisted Miura (G, \hbar) -opers and the set of nondegenerate polynomial solutions of the QQ -system:

$$\begin{aligned} \tilde{\xi}_i Q_-^i(\textcolor{brown}{u}) Q_+^i(\textcolor{blue}{\hbar u}) - \xi_i Q_-^i(\textcolor{blue}{\hbar u}) Q_+^i(\textcolor{brown}{u}) = \\ \Lambda_i(\textcolor{brown}{u}) \prod_{j>i} \left[Q_+^j(\textcolor{blue}{\hbar u}) \right]^{-a_{ji}} \prod_{j<i} \left[Q_+^j(\textcolor{brown}{u}) \right]^{-a_{ji}}, \quad i = 1, \dots, r, \end{aligned}$$

where $\tilde{\xi}_i = \zeta_i^{-1} \prod_{j>i} \zeta_j^{-a_{ji}}$, $\xi_i = \zeta_i \prod_{j<i} \zeta_j^{a_{ji}}$.

P. Koroteev, D. Sage, A.Z.,

(SL(N), q)-opers, the q-Langlands correspondence, and quantum/classical duality, Comm. Math. Phys. '21

E. Frenkel, P. Koroteev, D. Sage, A.Z.,

q-*Ops*, *QQ*-*Systems*, and *Bethe Ansatz*, to appear in *J. Eur. Math. Soc.*'24

Theorem.

There is a one-to-one correspondence between the set of nondegenerate Z -twisted Miura (G, \hbar) -opers and the set of nondegenerate polynomial solutions of the QQ-system:

$$\tilde{\xi}_i Q_-^i(u) Q_+^i(\hbar u) - \xi_i Q_-^i(\hbar u) Q_+^i(u) = \Lambda_i(u) \prod_{j>i} [Q_+^j(\hbar u)]^{-a_{ji}} \prod_{j<i} [Q_+^j(u)]^{-a_{ji}}, \quad i = 1, \dots, r,$$

where $\tilde{\xi}_i = \zeta_i^{-1} \prod_{j>i} \zeta_j^{-a_{ji}}$, $\xi_i = \zeta_i \prod_{j<i} \zeta_j^{a_{ji}}$.

P. Koroteev, D. Sage, A.Z.,

(SL(N), q)-opers, the q-Langlands correspondence, and quantum/classical duality, Comm. Math. Phys. '21

E. Frenkel, P. Koroteev, D. Sage, A.Z.,

q-Opers, QQ-Systems, and Bethe Ansatz, to appear in J. Eur. Math. Soc. '24

In ADE case this QQ-system correspond to the Bethe ansatz equations. Beyond simply-laced case: “folded integrable models”, based on $\widehat{\mathfrak{g}}$, see

E. Frenkel, D. Hernandez, N. Reshetikhin,

Folded quantum integrable models and deformed W-algebras, Lett. Math. Phys. '22

Introduction

Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

- **Cluster algebras** in Bethe ansatz through the notion of (G, \hbar) -Wronskians: QQ-systems via generalized minors of Fomin, Zelevinsky.

P. Koroteev, A.Z.,

q-Opers, QQ-systems, and Bethe Ansatz II: Generalized Minors, *Crelle J.*'23

- **Quantum-classical duality**: relation between classical multiparticle systems and spin chain systems through the natural coordinate change on (G, \hbar) -opers.

P. Koroteev, P. Pushkar, A. Smirnov, A.Z., *Quantum K-theory of Quiver Varieties and Many-Body Systems*, *Selecta Math.*'21;

P. Koroteev, D. Sage, A.Z., *$(SL(N), q)$ -opers, the q -Langlands correspondence, and quantum/classical duality*, *Commun. Math. Phys.* '21;

P. Koroteev, A.Z., *The Zoo of Opers and Dualities*, *IMRN*'24

- **3D mirror symmetry**: pairs of symplectic resolutions with similar properties. Coulomb and Higgs branches of 3D gauge theories. Enumerative geometry: $\{a_i\}$ - vs $\{z_i\}$ - qKZ equations.

P. Koroteev, A.Z., *Toroidal q -Opers*, *J. Inst. Math. Jussieu*'23

P. Koroteev, A. Z., *3d Mirror Symmetry for Instanton Moduli Spaces*, *Commun. Math. Phys.*'23

- Relation to **quantum q -Langlands correspondence**: (G, \hbar) -opers \rightarrow scalar differential operators for $q \rightarrow 1$ limit of conformal blocks for $W_{q,t}(L\mathfrak{g})$.

Introduction

Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

- ▶ **Cluster algebras** in Bethe ansatz through the notion of (G, \hbar) -Wronskians: QQ-systems via generalized minors of Fomin, Zelevinsky.

P. Koroteev, A.Z.,

q-Opers, QQ-systems, and Bethe Ansatz II: Generalized Minors, *Crelle J.*'23

- ▶ **Quantum-classical duality**: relation between classical multiparticle systems and spin chain systems through the natural coordinate change on (G, \hbar) -opers.

P. Koroteev, P. Pushkar, A. Smirnov, A.Z., *Quantum K-theory of Quiver Varieties and Many-Body Systems*,

Selecta Math.'21;

P. Koroteev, D. Sage, A.Z., *$(SL(N), q)$ -opers, the q -Langlands correspondence, and quantum/classical duality*,

Commun. Math. Phys. '21;

P. Koroteev, A.Z., *The Zoo of Opers and Dualities*, *IMRN*'24

- ▶ **3D mirror symmetry**: pairs of symplectic resolutions with similar properties. Coulomb and Higgs branches of 3D gauge theories. Enumerative geometry: $\{a_i\}$ - vs $\{z_i\}$ - qKZ equations.

P. Koroteev, A.Z., *Toroidal q -Opers*, *J. Inst. Math. Jussieu*'23

P. Koroteev, A. Z., *3d Mirror Symmetry for Instanton Moduli Spaces*, *Commun. Math. Phys.*'23

- ▶ Relation to **quantum q -Langlands correspondence**: (G, \hbar) -opers \rightarrow scalar differential operators for $q \rightarrow 1$ limit of conformal blocks for $W_{q,t}({}^L\mathfrak{g})$.

Introduction

Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

- ▶ **Cluster algebras** in Bethe ansatz through the notion of (G, \hbar) -Wronskians: QQ-systems via generalized minors of Fomin, Zelevinsky.

P. Koroteev, A.Z.,

q-Opers, QQ-systems, and Bethe Ansatz II: Generalized Minors, *Crelle J.*'23

- ▶ **Quantum-classical duality**: relation between classical multiparticle systems and spin chain systems through the natural coordinate change on (G, \hbar) -opers.

P. Koroteev, P. Pushkar, A. Smirnov, A.Z., *Quantum K-theory of Quiver Varieties and Many-Body Systems*,

Selecta Math.'21;

P. Koroteev, D. Sage, A.Z., *$(SL(N), q)$ -opers, the q -Langlands correspondence, and quantum/classical duality*,

Commun. Math. Phys. '21;

P. Koroteev, A.Z., *The Zoo of Opers and Dualities*, *IMRN*'24

- ▶ **3D mirror symmetry**: pairs of symplectic resolutions with similar properties. Coulomb and Higgs branches of 3D gauge theories. Enumerative geometry: $\{a_i\}$ - vs $\{z_i\}$ - qKZ equations.

P. Koroteev, A.Z., *Toroidal q -Opers*, *J. Inst. Math. Jussieu*'23

P. Koroteev, A. Z., *3d Mirror Symmetry for Instanton Moduli Spaces*, *Commun. Math. Phys.*'23

- ▶ Relation to **quantum q -Langlands correspondence**: (G, \hbar) -opers \rightarrow scalar differential operators for $q \rightarrow 1$ limit of conformal blocks for $W_{q,t}({}^L\mathfrak{g})$.

Introduction

Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

- ▶ **Cluster algebras** in Bethe ansatz through the notion of (G, \hbar) -Wronskians: QQ-systems via generalized minors of Fomin, Zelevinsky.

P. Koroteev, A.Z.,

q-Operators, QQ-systems, and Bethe Ansatz II: Generalized Minors, *Crelle J.*'23

- ▶ **Quantum-classical duality**: relation between classical multiparticle systems and spin chain systems through the natural coordinate change on (G, \hbar) -opers.

P. Koroteev, P. Pushkar, A. Smirnov, A.Z., *Quantum K-theory of Quiver Varieties and Many-Body Systems*, *Selecta Math.*'21;

P. Koroteev, D. Sage, A.Z., *$(SL(N), q)$ -opers, the q -Langlands correspondence, and quantum/classical duality*, *Commun. Math. Phys.* '21;

P. Koroteev, A.Z., *The Zoo of Opers and Dualities*, *IMRN*'24

- ▶ **3D mirror symmetry**: pairs of symplectic resolutions with similar properties. Coulomb and Higgs branches of 3D gauge theories. Enumerative geometry: $\{a_i\}$ - vs $\{z_i\}$ - qKZ equations.

P. Koroteev, A.Z., *Toroidal q -Opers*, *J. Inst. Math. Jussieu*'23

P. Koroteev, A. Z., *3d Mirror Symmetry for Instanton Moduli Spaces*, *Commun. Math. Phys.*'23

- ▶ Relation to **quantum q -Langlands correspondence**: (G, \hbar) -opers \rightarrow scalar differential operators for $q \rightarrow 1$ limit of conformal blocks for $W_{q,t}({}^L\mathfrak{g})$.

Introduction

Quantum Integrable models

Enumerative geometry and Bethe ansatz

QQ-systems and (G, \hbar) -opers

Applications

Thank you!